	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria	SEVERITY			
	Evaluation							
	SOIL							

Soil Erosion - Sheet and Rill	Detachment and transport of soil particles caused by rainfall splash and runoff degrade soil quality.	Sheet and rill erosion does not exceed the Soil Loss Tolerance "T".	Same as National	•	RUSLE2 (Primary) Visual assessment (pedestals, rills)	Severe = >4T Moderate = 2T to 4T Low = >T to 2T Visual assessment will be done to photo document concern (location, extent and severity).
Soil Erosion - Ephemeral Gully	Small channels caused by surface water runoff degrade soil quality and tend to increase in size. On cropland, they can be obscured by heavy tillage.	Surface water runoff is controlled sufficiently to stabilize the small channels and prevent reoccurrence of new channels.	Same as National	•	Volume calculation (Primary) Visual assessment	Severe = Soil loss increased > 3T Moderate = Soil loss increased 1T to 3T Low Soil loss increased <1T Visual assessment will be done to photo document concern (location, extent and severity).
Soil Erosion - Classic Gully	Deep, permanent channels caused by the convergence of surface runoff degrade soil quality. They enlarge progressively by headcutting and lateral widening.	Surface water runoff is controlled sufficiently to stop progression of headcutting and widening.	Same as National	•	Volume calculation (Primary) Visual assessment Aerial photo trend analysis	Severe = Soil loss increased > 3T Moderate = Soil loss increased 1T to 3T Low Soil loss increased <1T Visual assessment will be done to photo document concern (location, extent and severity). Aerial photo trend analysis will be done to document time of erosion and use for baseline for monitoring success.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria	SEVERITY			
	Evaluation							
	SOIL							

Soil Erosion - Streambank	Accelerated loss of streambank soils restricts land and water use and management.	Accelerated streambank soil loss does not exceed a level commensurate with upstream land use and normal geomorphological processes on site.	Same as National	•	Volume calculation (Primary) Aerial photo trend analysis Visual assessment, e.g., Stream Visual Assessment Protocol, (Secondary)	Severe = Soil loss > 1/4T / lin ft. Moderate = Soil loss between 1/8 to 1/4T / lin. Ft. Low Soil loss < 1/8 T/ lin. Ft. Aerial photo trend analysis will be done to document time of erosion and use for baseline for monitoring success. Visual assessment will be done using the SVAP and photo documented to record the concern (location, extent and severity).
Soil Condition - Organic Matter Depletion	Soil organic matter has or will diminish to a level that degrades soil quality.	Soil Conditioning Index is positive.	Same as National	•	Soil Conditioning Index (Primary)	Severe = > - 0.30 Moderate = between - 0.10 and - 0.30 Low = 0.0 to - 0.10
Soil Condition - Compaction	Compressed soil particles and aggregates caused by mechanical compaction adversely affect plant-soilmoisture relationships.	Mechanically compacted soils are renovated sufficiently to restore plant root growth and/or water movement.	Same as National	•	Dial penetrometer (Primary)	Severe = > 300 Moderate = between 200 and 300 Low = 0 to 200

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria Evaluation	SEVERITY			
	SOIL							

Soil Condition - Contaminants - Animal Waste and Other Organics	Nutrient levels from applied animal waste and other organics restrict desired use of the land.	Nutrient application levels do not exceed soil storage/plant uptake capacities based on soil test recommendations and risk analysis results.	Same as National	•	Rhode Island Phosphorus Index (Primary) Soil test (Secondary)	Severe = RIPI 65-75 Moderate = RIPI 39-64 Low= RIPI 25-38 Soil test results are needed to identify the limiting nutrient and assess recommendations for applications and establish a base line.
Soil Condition Contaminants Commercial Fertilizer	Over application of nutrients degrades plant health and vigor, or exceeds the soil capacity to retain nutrients.	Soil nutrient levels do not exceed crop needs based on realistic yield goals and appropriate pH levels are maintained.	Same as National	•	Rhode Island Phosphorus Index (Primary) Soil test (Secondary)	Severe = RIPI 65-75 Moderate = RIPI 39-64 Low= RIPI 25-38 Soil test results are needed to identify the limiting nutrient and assess recommendations for applications and establish a base line.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP						
Natural Resource	Description of Concern	National Quality	State Quality	Assessment Tools for			
Concern		Criteria	Criteria	Quality Criteria Evaluation			
	WATER						

Water Quantity - Excessive Seepage	Subsurface water oozing to the surface restricts land use and management.	Subsurface water is managed to limit periods of saturation that are unfavorable to the present or intended land use. Management complies with wetland policies.	Same as National	•	Visual Assessment (physical presence of water, prevalence of hydrophytic vegetation, etc.) (Primary)	Visual assessment will be done to photo document concern (location, extent and severity). On cropland, orchards, vineyards: Severe = large area of concern that occupies > 10% of the area, prohibits access to and/or management on the area for greater then 30 days during the growing season (April 1 to August 30). Moderate = areas where 2 to 10% of the site are in accessible and/or management on for two to four weeks of the growing season. Low = small, incidental areas (less then 2%) where seepage limits access to and/or management on area for short durations of the growing season (typically, immediately after rainfall events – less
				•	Client interview and area measurements (Secondary)	then a two week period). Client interview is to establish history of concern (how long seepage has been occurring, time of year, duration and volume estimates). This information should support the claim of restricted use and management of site.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria				
	Evaluation							
	WATER							

Water Quantity -	The land becomes inundated restricting	Excess water amounts and/or rates		•	Visual assessment	Visual assessment will be done to photo document concern (location, extent and
Excessive	land use and	of flow are controlled	Same as		(Primary)	severity).
Runoff,	management.	consistent with	National		01 \1"	Covers - erece (> 100/ of the erec)
Flooding, or Ponding		desired present or intended land use goals and wetland policies.		•	Stream Visual Assessment Protocol	Severe = areas (> 10% of the area) affected by runoff, flooding or ponding that limits access to and/or management on the area for greater then 30 days during
		policies.		•	Hydrologic	the growing season (April 1 to August 30).
					models, e.g. HECRAS,TR- 20,TR-55	Moderate = areas (2 to 10% of the area) affected by runoff, flooding or ponding that limits access to and/or management on the area for two to four weeks of the growing season.
						Low = small, incidental areas (less then 2%) affected by runoff, flooding or ponding that limits access to and/or management on area for short durations of the growing season (typically, immediately after rainfall events – less then a two week period).
				•	Client interview (Secondary)	Client interview is to establish history of concern (how long runoff, flooding or ponding has been occurring, time of year, duration and volume estimates). This information should support the claim of restricted use and management of site.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria Evaluation				
	WATER							

Water Quantity -	Water saturates upper soil layers restricting	Subsurface water is managed to limit		•	Visual assessment of	Visual assessment will be done to photo document concern (location, extent and
Excessive Subsurface	land use and management.	periods of saturation compatible with the	Same as National		soil cores and coring holes	severity).
Water		present or intended land use and wetland			(Primary)	On cropland, orchards, vineyards:
		policies.		•	Plant quality and quantity measurements	Severe = large area of concern that occupies > 10% of the area, prohibits access to and/or management on the area for greater then 30 days during the growing season (April 1 to August 30).
						Moderate = areas where 2 to 10% of the site are in accessible and/or management on for two to four weeks of the growing season.
						Low = small, incidental areas (less then 2%) where seepage limits access to and/or management on area for short durations of the growing season (typically, immediately after rainfall events – less then a two week period).
				•	Client interview (Secondary)	Client interview is to establish history of concern (how long excessive subsurface water has been occurring, time of year, duration and volume estimates). This information should support the claim of restricted use and management of site.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural	Description of	National	State	Assessment Tools						
Resource	Concern	Quality	Quality	for						
Concern		Criteria	Criteria	Quality Criteria						
				Evaluation						
	WATER									

Water Quantity - Inefficient Water Use on Irrigated Land	Limited water supplies are not optimally utilized.	Land and water management is planned and coordinated to provide optimal use of natural and applied moisture.	Same as National	Farm Irrigation Rating Index and/or Table below (Primary System Type Trickle or Drip Irrigation Point Source Emitter Spray Emitter Continuous Tape		Severe = >20% improvement in system efficiency Moderate = 10-20% improvement in system efficiency Low = 0-10% improvement in system efficiency, and/or no IWM in place for high-volume system.
				Sprinkler Handline and Wheel Big Gun (stationary) Traveling Gun Solid Set Center Pivot w/overhead spray Center Pivot w/drop down emitters Linear Move w/overhead spray Linear Move w/drop down emitters When converting from one system to another, the difference in system efficiency = increase/improvement in system efficiency for adding buried mainline. IWM is required for all high-volume systems. If not, rank as low and should be an incentive in ranking. Visual assessment (Second	70 60 67 85 85 90 85 90	Visual assessment will be done to photo document concern (location, extent and severity).

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural	Description of	National	State	Assessment Tools						
Resource	Concern	Quality	Quality	for						
Concern		Criteria	Criteria	Quality Criteria						
				Evaluation						
	WATER									

Water Quantity – Insufficient Flows in Water Courses	Water flows are not consistently available in sufficient quantities to support ecological processes and land use and management.	Authorized uses and management of water are coordinated to minimize the impacts on water course flows.	Same as National	•	Gauge Station data / Water flow records (Primary) Pumping records Hydrologic models, if available Fish and Wildlife Survey data Visual assessment (Secondary)	Based upon expected low flows as determined by USGS Low Flow Report, Water Availability studies, and other USGS reports. Severe = Stream withdrawals creating low flow watersheds, including Beaver, Chipuxet, Blackstone, Hunt and Nipmuc rivers. Moderate = Stream withdrawals creating low flows in the Wood/Pawcatuck watershed (except for Chipuxet and Beaver listed above), Ten Mile, Branch, Jamestown, Aquidneck Island and Bristol County drinking water supply areas, and withdrawals in all cold water fisheries waters (refer to draft list at RIDEM Office of Water Resources – Alisa Richardson). Low = All other watersheds Visual assessment will be done to photo document concern (location, extent and severity).
Water Quality - Harmful Levels of Pesticides in Groundwater	Residues resulting from the use of pest control chemicals degrade groundwater quality.	Pesticides are applied, stored, handled, disposed of, and managed so that groundwater uses are not adversely affected	Same as National	•	WIN-PST (Windows Pesticide Screening Tool – USDA/NRCS) (Primary)	Severe = High or Very High* Moderate = Intermediate* Low = Very Low or Low* *WIN-PST Rating for leaching

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural Resource	e Concern Quality	National Quality	State Quality	Assessment Tools for						
Concern		Criteria	Criteria	Quality Criteria Evaluation						
	WATER									

Water Quality - Excessive Nutrients and Organics in Groundwater	Pollution from natural or human induced nutrients such as N, P, and organics (including animal and other wastes) degrades groundwater quality.	Nutrients and organics are stored, handled, disposed of, and applied such that groundwater uses are not adversely affected.	Same as National	•	Rhode Island Phosphorus Leaching Index (RIPI) (Primary) Nitrate Leaching Index Severity		Distance be (Sole Sou wetland watercour Slope of Contributing Area	RIPI 39-25-38 areas what r 45 days not be grea): etween HU rce Aquifer i, third , sese) without Soil Texture Class I < 200'	A and Sens public well cond or first an adequated Soil Texture Class II	a year and sed within the sitive Area , state id order e buffer. Soil Texture Class III < 300'
Water Quality - Harmful Levels of Petroleum in Groundwater	Fuel, oil, gasoline and other hydrocarbons present in toxic amounts degrade groundwater quality.	Petroleum products are used, stored, handled, disposed of, and managed such that groundwater uses are not adversely affected.	Same as National	•		Soil Texture Soil Texture I water e maps y)	3-5% 0-3% Class I - sands Class III - all o Severe = P Moderate = 2.0 in./hr. Low = Pern Visual assed document of severity).	y clay loan ther texturate remeabil remeability.	is and sand all classes ity > 2.0 i bility betv < 0.2 in./h will be doi	n./hr. veen 0.2 and r. ne to photo

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural	Description of	National	State	Assessment Tools						
Resource	Concern	Quality	Quality	for						
Concern		Criteria	Criteria	Quality Criteria						
				Evaluation						
	WATER									

Water Quality	Pest control	Pesticides are		•	WIN-PST	Severe = High or Very High*
- Harmful	chemicals present in	applied, stored,			(Windows	Moderate = Intermediate*
Levels of	toxic amounts	handled, disposed of,			Pesticide	Low = Very Low or Low*
Pesticides in	degrade surface	and managed such	Same as		Screening Tool	
Surface Water	water quality.	that surface water	National		- USDA/NRCS)	*WIN-PST Rating for leaching
		uses are not				_
		adversely affected				

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural	Description of	National	State	Assessment Tools						
Resource	Concern	Quality	Quality	for						
Concern		Criteria	Criteria	Quality Criteria						
				Evaluation						
	WATER									

Water Quality - Excessive Nutrients and Organics in Surface Water	Pollution from natural or human induced nutrients such as N, P, and organics (Including animal and other wastes) degrades surface water quality.	Nutrients and organics are stored, handled, disposed of, and managed such that surface water uses are not adversely affected.	Same as National	F	Rhode Island Phosphorous index (RIPI) (Primary)		For Cropland, Hayland and Pasture: Severe = RIPI 65-75 Moderate = RIPI 39-64 Low= RIPI 25-38 For HUAs (areas where livestock are confined for 45 days or more a year and forage will not be grown for feed within the confined area):			
				\	Visual Assess Protoco	Soil Textur Soil Textur Stream ment	(Sole Sour	rce Aquifer, third, se se) without Soil Texture Class I < 200' < 150' < 100' s and loam y clay loam ther textura	ns and sandy al classes establish	state id order e buffer. Soil Texture Class III < 300' < 250' < 200'

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural Resource	e Concern Quality	National Quality	State Quality	Assessment Tools for						
Concern		Criteria	Criteria	Quality Criteria Evaluation						
	WATER									

Water Quality - Excessive Suspended Sediment and Turbidity in Surface Water	Pollution from mineral or organic particles degrades surface water quality.	Movement of mineral and organic particles is managed such that surface water uses are not adversely affected.	Same as National	•	SVAP (Stream Visual Assessment Protocol – USDA/NRCS) (Primary)	SVAP will be used as the primary tool and to establish a baseline for use in future assessments. Severe = Moderate = Low =
				•	Visual assessment	Visual assessment will be done to photo document concern (location, extent and severity).
				•	Client interview	Client interview is to establish history of concern (how long excessive suspended sediment has been occurring, time of year and duration).
Water Quality - Harmful Temperatures of Surface Water	Undesired thermal conditions degrade surface water quality.	Use and management of land and water are coordinated to minimize impacts on surface water temperatures.	Same as National	•	SVAP (Stream Visual Assessment Protocol – USDA/NRCS) – canopy cover (Primary) and/or Thermal pollution identified as a limiting factor in state collected data bases.	SVAP will be used as the primary tool and to establish a baseline for use in future assessments. Severe = Moderate = Low =

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria				
				Evaluation				
	WATER							

Water Quality - Harmful Levels of Pathogens in Surface Water	Kinds and numbers of viruses, protozoa, and bacteria are present at a level that degrades surface water quality.	Materials that harbor pathogens are stored, handled, disposed of, applied, and managed such that surface water uses are not adversely affected.	Same as National	•	State 303d listed	Severe = Primary pollutant of concern on impacted waterbody Moderate = Identified as a possible pollutant of concern on impacted waterbody. Low = Data from other sources indicating occurrence of pathogens within the watershed adjacent to a listed waterbody.
Water Quality - Harmful Levels of Petroleum in Surface Water	Fuel, oil, gasoline and other hydrocarbons present in toxic amounts degrade surface water quality.	Petroleum products are used, stored, handled, and disposed of such that groundwater uses are not adversely affected.	Same as National	•	State 303d listed	Severe = Primary pollutant of concern on impacted waterbody Moderate = Identified as a possible pollutant of concern on impacted waterbody. Low = Data from other sources indicating occurrence of pathogens within the watershed adjacent to a listed waterbody.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria				
				Evaluation				
	AIR							

Air Quality – Land use and management operations produced by the control of the control operation of the control operation ope	Odor-producing		•	Olfactory	Assessment will be based upon prevailing
Airborne Odors Objectionable of from such sour confined liveste animal waste, waste storage waste lagoons, application of a waste, and oth organics.	to mitigate potential nuisance impacts and meets all applicable Tribal, State, and Local regulations. areas, field nimal	Same as National		assessment (include client interview and statements from regulatory or administrative offices of local municipalities) (Primary) and use the Agricultural Waste Management Field Handbook (AWMFH) (Secondary)	winds and landscape elements such as building arrangement, landform, and vegetation from which odors may emanate and visual resource concerns may compound the concern. Severe = Source of odor is within 100 feet of property line, neighboring property or other targeted concern. Moderate = Source of odor is between 100 and 200 feet of property line, neighboring property or other targeted concern. Low = Source of odor is greater than 200 feet of property line, neighboring property or other targeted concern.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural	Description of	National	State	Assessment Tools				
Resource	Concern	Quality	Quality	for				
Concern		Criteria	Criteria	Quality Criteria				
				Evaluation				
	PLANTS							

<u> </u>	ID: () (1		
Plant – Condition- Productivity, Health and Vigor	Plants do not produce the yields, quality, and soil cover to meet client objectives.	Selected plants on or planned for the site are sufficiently productive to meet or exceed client needs. Cropland: A healthy stand with vigorous growth produces at least 75% of site potential.	Same as	 Client interview (Primary) For cropland and hayland. 	Client interview is to establish history of plant yield (records to include amounts and dates) to be compared to targeted yields listed in the Yield Tables in Section II A. of the RI eFOTG. Severe if Yield < 50% of target Moderate if Yield between 50 to 75% of target Low if Yield at 75% or better of target
		Pastureland: Forage yields are at least 75% of high management estimates cited yields listed in the Yield Tables in Section II A. of the RI eFOTG. Hayland: Forage Yields at least 75% of high mgt. estimates cited yields listed in the Yield Tables in Section II A. of the RI eFOTG. Forestland/ Agroforest: Forests consist of healthy stands with vigorous growth having a stand density within 25% of optimum stocking on a stems/acre basis.		 For Pasture Use The RI Pasture Condition Score Sheet Indicators: percent desirable plants, plant cover, plant residue, plant vigor, percent legumes and uniformity of use. For Forestland, use the forest stewardship plan data submitted by a consulting forester. 	Severe if PCS is 2 or less Moderate if PCS >2 and less then 4 Low if PCS sub total > 4 Severe if Yield < 50% of target Moderate if Yield between 50 to 75% of target Low if Yield at 75% or better of target
					Page 15 of 15

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria Evaluation				
				Evaluation				
	PLANTS							

Plant Condition - Threatened or Endangered Plant Species	Plant populations and /or habitat quantity and quality have reached a level that one or more plant species are in danger of or threatened with extinction.	Threatened and endangered plant species and/or habitats they occupy are managed to avoid actions that would reduce their current population, health, or sustainability.	Same as National	Consultation with appropriate federal, state, and local agencies/groups (Primary) Client interviews	Severe = T&E species identified within CTU Moderate = T&E species identified within range of CTU for possible impact by management for specie. Low = T&E species not present but CTU has been identified as a potential site by specialist. Client interview is to establish know history/incidence/management of T&E on site.
Plant Condition - Noxious and Invasive Plants	The site has noxious or invasive plants present.	The site is managed to control noxious and invasive plants and to minimize their spread.	Same as National	Inventory site using RI HAP for Invasive Plants (Primary) Client interviews	Severe if HIS = 0.1 but less then 0.5 Moderate if HIS = 0.5 but less then 1.0 Low if HIS is = 1.0 Client interview is to establish know history/incidence/management of noxious or invasive plants on site.

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural	Description of	National	State	Assessment Tools				
Resource	Concern	Quality	Quality	for				
Concern		Criteria	Criteria	Quality Criteria				
				Evaluation				
	ANIMALS							

Fish and Wildlife - Inadequate Food	Quantity and quality of food is unavailable to meet the life history requirements of the species or guild of species of concern	Food availability meets the life history requirements of the species or guild of species of concern.	Same as National	 Inventory site using RI HAP for Invasive Plants (Primary) Visual assessment Visual severe if HIS = 0.1 but less then 0.5 Moderate if HIS = 0.5 but less then 1.0 Visual assessment will be done to photo document concern (location, extent and severity).
Fish and Wildlife – Inadequate Cover/Shelter	Cover/shelter for the species of concern is unavailable or inadequate. For aquatic species, this includes lack of hiding, thermal, and/or refuge cover	The ecosystem or habit types support the necessary plant species in the kinds, amounts, and physical structure; and the connectivity of fish and wildlife cover is adequate to support, over time, the species of concern.	Same as National	 Inventory site using RI HAP for Invasive Plants (Primary) Visual assessment Visual severity) Visual assessment will be done to photo document concern (location, extent and severity).
Fish and Wildlife – Inadequate Water	The quantity and quality of water is unacceptable for the species of concern	The quantity and quality of water meets the life history requirements of the species of concern.	Same as National	 Inventory site using RI HAP for Invasive Plants (Primary) Visual assessment Visual severity) Visual assessment will be done to photo document concern (location, extent and severity).

	Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP							
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria				
Concern		Officeria	Officeria	Evaluation				
	ANIMALS							

Fish and Wildlife - Threatened and Endangered Species	Fish and wildlife populations and/or habitat quantity and quality have reached a level that one or more species are in danger of or threatened with extinction.	Threatened and endangered fish and wildlife species and/or habitats they occupy are managed to avoid actions that would reduce their current population, health, or sustainability.	Same as National	Consultation with appropriate federal, state, and local agencies /groups (Primary) Client interviews	Severe = T&E species identified within CTU Moderate = T&E species identified within range of CTU for possible impact by management for specie. Low = T&E species not present but CTU has been identified as a potential site by specialist. Client interview is to establish know history/incidence/management of T&E on site.
Domestic Animals – Inadequate Quantities and Quality of Feed and Forage	Total feed and forage is insufficient to meet the nutritional and production needs of the kinds and classes of livestock	Feed and forage including supplemental nutritional requirements are provided to meet production goals for the kinds and classes of livestock. Native grazers are factored into the total feed and forage balance computations.	Same as National	 Measured inventory Visual assessment 	Severe = Moderate = Low = Visual assessment will be done to photo document concern (location, extent and severity).

Rhode Island NRCS FY 2005 Resource Concerns and Quality Criteria for EQIP									
Natural Resource Concern	Description of Concern	National Quality Criteria	State Quality Criteria	Assessment Tools for Quality Criteria					
Concern		Officeria	Officeria	Evaluation					
ANIMALS									

Domestic Animals – Inadequate Shelter	Livestock are not protected sufficiently to meet the production goals for the kinds and classes of livestock	Artificial and/or natural shelter is provided to meet production goals for the kinds and classes of livestock.	Same as National	 Inventory of facilities and their capacities (Primary) Visual assessment Aerial photo analysis 	If existing percentage of the targeted* paddock or HUA is: Severe = < 50% Moderate = between 50 – 80% Low = > 80 but under 100% *Target horse paddock area is 180 sq. ft. per animal and target cattle HUA area is 100 sq. ft. per animal. Visual assessment will be done to photo document concern (location, extent and severity). Paddock sizes must be sufficient for animals in sacrifice areas or HUAs.
Domestic Animals – Inadequate Stock Water	The quantity, quality and distribution of drinking water is insufficient to meet the production goals for the kinds and classes of livestock	Sufficient water of acceptable quality is provided and adequately distributed to meet production goals for the kinds and classes of livestock. To reduce potential for water contamination, watering facilities are constructed or modified to minimize mortality to indigenous wildlife.	Same as National	 Inventory of distribution needs (Primary) Visual assessment Aerial photo analysis 	Severe = Moderate = Low = Visual assessment will be done to photo document concern (location, extent and severity).